

Shelf Life

EPIKURE[™] Curing Agent 3233

This product has a minimum shelf life (no maximum) of three (3) years from date of certification at recommended storage conditions.

The recommended storage conditions for this product is 75 ± 25°F max., dry and free of contamination.

"Shelf Life" is the average time during which product properties can be expected to remain within product specifications set out in the applicable sales contract under appropriate storage and handling conditions, and assuming there has been no contamination. As an accommodation to our customers, we can provide information as to the typical MINIMUM Shelf Life of our products; however, we do not warrant Shelf Life nor assume any responsibility for such information. After expiration of the MINIMUM Shelf Life, or in situations where the product may not have been handled or stored properly or where contamination may have occurred, the customer is advised to evaluate the product for conformity with specifications and suitability for the use intended. Although there are exceptions, in many cases, if properly handled and stored, product will remain usable for its normal applications for a significant period beyond the MINIMUM Shelf Life.

The date of certification for the product is reported on the Certificate of Analysis. Shelf life is based on the Certification Date. The manufacturing date can be determined from the Lot Number as follows:

- For Products Manufactured in the 2nd half of 2007 and beyond - The first number (position 3) in the lot number sequence represent the year (1 = 2011, 2 = 2012, and so on, with 0 = 2010) The next letter in the lot number sequence represents the month (A = January, B = February, and so on, with L = December). For example - Lot Number is LL1G0001, indicates the product was manufactured in July 2011.

To preserve product quality and prevent discoloration, it is recommended that a nitrogen blanket be maintained on the headspace of an opened container and that storage / handling temperatures in excess of 50°C (122°F) be avoided. Spillage around the opening of the container from dispensing operations will form a crystalline residue. This residue is not soluble in the curing agent nor the resin, and certain measures should be taken to prevent it from contaminating the remaining contents of the container. The crystalline residue should be removed with a warm damp wash cloth PRIOR to re-opening the container for dispensing.

Our standard sales contracts typically provide a warranty by us that our products will, upon delivery to the customer at the contract delivery point(s), comply with all product specifications set out in the contract. However, once delivery has occurred, since we no longer have possession or control over the product, including its handling and storage, we cannot continue to be responsible for product quality. (Please see Warranty Disclaimer)

DISCLAIMER

The information provided herein was believed by Hexion Inc. ("Hexion") to be accurate at the time of preparation or prepared from sources believed to be reliable, but it is the responsibility of the user to investigate and understand other pertinent sources of information, to comply with all laws and procedures applicable to the safe handling and use of the product and to determine the suitability of the product for its intended use. All products supplied by Hexion are subject to Hexion's terms and conditions of sale. **HEXION MAKES NO WARRANTY, EXPRESS OR IMPLIED, CONCERNING THE PRODUCT OR THE MERCHANTABILITY OR FITNESS THEREOF FOR ANY PURPOSE OR CONCERNING THE ACCURACY OF ANY INFORMATION PROVIDED BY HEXION**, except that the product shall conform to Hexion's specifications. Nothing contained herein constitutes an offer for the sale of any product.